

PARADICHLOROBENZENE

fact sheet

What is paradichlorobenzene?

Paradichlorobenzene is used as a fumigant insecticide to control clothes moths. It is also found in deodorant blocks made for trash cans and toilets. Paradichlorobenzene was first registered for use in the United States in 1942, and it is sometimes called 1,4-dichlorobenzene.

What are some products that contain paradichlorobenzene?

Mothballs containing paradichlorobenzene are solids that turn into toxic gas that kills moths. In 2010, there are over 30 products registered with the U.S. Environmental Protection Agency that contain paradichlorobenzene.

How does paradichlorobenzene work?

The vapor of paradichlorobenzene is toxic to insects. In humans and other animals, paradichlorobenzene is broken down in the body to form other compounds that may be harmful to cells or organs such as the liver.



clothes moth, photo credit:
Patrick Clement, flickr

IMPORANT: Always *follow label instructions* and take steps to *avoid exposure*. If any exposures occur, be sure to follow the First Aid instructions on the product label carefully. For additional treatment advice, contact the Poison Control Center at 800-222-1222. If you wish to discuss a *pesticide problem*, please call 1-800-858-7378.

How might I be exposed to paradichlorobenzene?


You can be exposed to a pesticide if you breathe it in, get it on your skin, or if you accidentally eat or drink something containing a pesticide. This can happen if you get some on your hands and don't wash them before eating or smoking. People are most likely to be exposed to paradichlorobenzene by breathing in the vapors. When you smell mothballs, you are inhaling the pesticide. Small children and pets are at risk of eating mothballs, because they look like candy or other treats.

What are some signs and symptoms from a brief exposure to paradichlorobenzene?

People who have been exposed to paradichlorobenzene have experienced nausea, vomiting, dizziness, fatigue, and headaches. Paradichlorobenzene vapor can also irritate the eyes and nasal passages. If paradichlorobenzene contacts the skin for a prolonged period, it can cause a burning sensation. If a pet eats a mothball made of paradichlorobenzene, they may have vomiting, tremors, and/or abdominal pain. Paradichlorobenzene may also cause kidney and liver damage in pets.

PARADICHLOROBENZENE

fact sheet



What happens to paradichlorobenzene when it enters the body?

In humans, paradichlorobenzene is distributed in the blood, fat, and breast milk. It is broken down into several other chemicals by the body and excreted in urine. Human volunteers who inhaled paradichlorobenzene exhaled half the dose. The amount of paradichlorobenzene in their blood dropped by more than 50% one hour after the exposure stopped.

In animals, paradichlorobenzene is rapidly absorbed through the lungs or gut, but more slowly through the skin. Paradichlorobenzene was found in the fat, liver, and kidneys. Smaller amounts were found in the blood plasma, lungs, and muscle. Paradichlorobenzene was eliminated from the body soon after the exposure stopped. When animals were exposed for long periods of time, their bodies began to break down the paradichlorobenzene faster, and tissue levels declined.

Is paradichlorobenzene likely to contribute to the development of cancer?

The World Health Organization (WHO) considered paradichlorobenzene possibly carcinogenic to humans based on studies with mice. The way paradichlorobenzene caused cancer in mice could possibly occur in humans as well. The U.S. EPA has classified it as “not likely to be carcinogenic to humans”.

Has anyone studied non-cancer effects from long-term exposure to paradichlorobenzene?

Rats and rabbits exposed to very high doses of paradichlorobenzene vapor while pregnant gained less weight than control animals. In another study, mother rats had fewer surviving young. No information was found regarding paradichlorobenzene and asthma or other chronic diseases.

Are children more sensitive to paradichlorobenzene than adults?

While children may be especially sensitive to pesticides compared to adults, there are currently no data to suggest that children have increased sensitivity specifically to paradichlorobenzene. Young children may be more at risk of eating mothballs because the mothballs may look like candy.

What happens to paradichlorobenzene in the environment?

Most of the paradichlorobenzene that gets into the environment will turn into vapor. It can also be broken down by bacteria or become attached to sediments in water. Paradichlorobenzene that binds to soil may be taken up by plants, and plant leaves may absorb paradichlorobenzene from the air. Paradichlorobenzene in air is broken down slowly by other chemicals. It has been found in rainwater and snow. Paradichlorobenzene has been found in groundwater close to a source of contamination. In air, its half-life is about 31 days.


NPIC is a cooperative agreement between Oregon State University and the U.S. Environmental Protection Agency (U.S. EPA). The information in this publication does not in any way replace or supercede the restrictions, precautions, directions, or other information on the pesticide label or any other regulatory requirements, nor does it necessarily reflect the position of the U.S. EPA.



Oregon State
University

PARADICHLOROBENZENE

fact sheet



Can paradichlorobenzene affect birds, fish, or other wildlife?

When researchers fed 10 ducks a diet containing 0.5% paradichlorobenzene for 35 days, three ducks died and the rest did not grow as well. Paradichlorobenzene is moderate to low in toxicity to fish, with differences in sensitivity by species. No information was found for the effects of paradichlorobenzene on bees.

Where can I get more information?

For more detailed information about paradichlorobenzene please visit the list of referenced resources below, call NPIC between 8:00 AM and 12:00 PM Pacific Time (11:00 AM to 3:00 PM Eastern Time), Monday - Friday, at 800-858-7378, or visit us on the web at npic.orst.edu. NPIC provides objective, science-based answers to questions about pesticides.

Date Reviewed: MM YYYY


Cite as: Gervais, J.; Luukinen, B.; Buhl, K.; Stone, D. 2010. *Paradichlorobenzene General Fact Sheet*; National Pesticide Information Center, Oregon State University Extension Services. npic.orst.edu/factsheets/PDBgen.html.

References:

1. Revised Reregistration Eligibility Decision (RED) Para-dichlorobenzene, EPA 738-R-07-010; U.S. Environmental Protection Agency, Office of Prevention, Pesticides, and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 2008.
2. Toxicological profile for dichlorobenzenes; U.S. Department of Health and Human Services, Agency for Toxic Substances and Disease Registry: Atlanta, GA, 2006.
3. Para-Dichlorobenzene: HED chapter of the Reregistration Eligibility Decision Document (RED); U.S. Environmental Protection Agency, Office of Prevention, Pesticides, and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office : Washington, DC, 2007.
4. Scuderi, R. Determination of para-dichlorobenzene releases from selected consumer products. Unpublished report, 1986, submitted to U.S Department of Health and Human Services prepared by Midwest Research Institute, Kansas City, MO. Toxicological profile for dichlorobenzenes; U.S. Department of Health and Human Services, Agency for Toxic Substances and Disease Registry: Atlanta, GA, 2006.
5. Hollingsworth, R. L.; Rowe, V. K.; Hoyle, H. R.; Spencer, H. C. Toxicity of paradichlorobenzene. *AMA Arch. Ind. Health* 1956, 14 (2), 138-147.
6. Lide, D. *CRC Handbook of Chemistry and Physics*; CRC Press Inc.: Boca Raton, FL, 1994; p 16-25.
7. Yan, R. M.; Chiung, Y. M.; Pan, C. Y.; Liu, J. H.; Liu, P. S. Effects of dichlorobenzene on acetylcholine receptors in human neuroblastoma SH-SY5Y cells. *Toxicol.* 2008, 253 (1-3), 28-35.
8. Gaines, T. B.; Linder, R. E. Acute toxicity of pesticides in adult and weanling rats. *Fundam. Appl. Toxicol.* 1986, 7, 299-308.

PARADICHLOROBENZENE


fact sheet



9. Morris, T. Primary eye irritation study in rabbits: para-dichlorobenzene. Unpublished lab project no. 91-8305-21(B), 1992, submitted to U.S. Environmental Protection Agency prepared by Hill Top Biolabs, Inc. Revised Reregistration Eligibility Decision (RED) Para-dichlorobenzene, EPA 738-R-07-010; U.S. Environmental Protection Agency, Office of Prevention, Pesticides, and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 2008.
10. Morris, T. Primary skin irritation study in rabbits: para-dichlorobenzene. Unpublished lab project no. 91-8305-21(A), 1992, submitted to U.S. Environmental Protection Agency prepared by Hill Top Biolabs, Inc. Revised Reregistration Eligibility Decision (RED) Para-dichlorobenzene, EPA 738-R-07-010; U.S. Environmental Protection Agency, Office of Prevention, Pesticides, and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 2008.
11. Morris, T. Delayed contact hypersensitivity in guinea pigs (Buehler technique): para-dichlorobenzene. Unpublished lab project no. 9-8305-21(C), 1992, submitted to U.S. Environmental Protection Agency prepared by Hill Top Biolabs, Inc. Revised Reregistration Eligibility Decision (RED) Para-dichlorobenzene, EPA 738-R-07-010; U.S. Environmental Protection Agency, Office of Prevention, Pesticides, and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 2008.
12. Newton, P. An acute inhalation toxicity study of para-dichlorobenzene in the rat. Unpublished lab project no. 89-8230, 1990, submitted to U.S. Environmental Protection Agency prepared by Bio/dynamics, Inc. Revised Reregistration Eligibility Decision (RED) Para-dichlorobenzene, EPA 738-R-07-010; U.S. Environmental Protection Agency, Office of Prevention, Pesticides, and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 2008.
13. Oehme, F. W.; Kore, A. M. Miscellaneous indoor toxicants. *Small Animal Toxicology*, 2nd ed.; Peterson, M. E.; Talcott, P. A. Eds.; Elsevier: St. Louis, MO, 2006; p 240.
14. DeClementi, C. Moth repellent toxicosis. *Vet. Med.* 2005, 24-28.
15. LaBonde, J. Toxicity in pet avian patients. *Semin. Avian Exot. Pet Med.* 1995, 4 (1), 23-31.
16. Reigart, J. R.; Roberts, J. R. Fumigants. *Recognition and Management of Pesticide Poisonings*, 5th ed.; U.S. Environmental Protection Agency, Office of Prevention, Pesticides, and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 1999; p 160.
17. Cotter, L. H. Paradichlorobenzene poisoning from insecticides. *N. Y. State J. Med.* 1953, 53 (14), 1690-1692.
18. Miyai, I.; Hirono, N.; Fujita, M.; Kameyama, M. Reversible ataxia following chronic exposure to paradichlorobenzene. *J. Neurol. Neurosurg. Psychiatr.* 1988, 51, 453-454.
19. Campbell, D. M.; Davidson, R. J. L. Toxic haemolytic anaemia in pregnancy due to a pica for paradichlorobenzene. *J. Obstet. Gynaecol. Br. Commonw.* 1970, 77, 657-659.
20. Hallowell, M. Acute haemolytic anemia following the ingestion of para-dichlorobenzene. *Arch. Dis. Child.* 1959, 34, 74-75.
21. Nalbandian, R. M.; Pearce, J. F. Allergic purpura induced by exposure to p-dichlorobenzene. *J. Am. Med. Assoc.* 1965, 194 (7), 828-829.
22. Aiso, S.; Arito, H.; Nishizawa, T.; Nagano, K.; Yamamoto, S.; Matsushima, T. Thirteen-week inhalation toxicity of p-dichlorobenzene in mice and rats. *J. Occup. Health* 2005, 47, 249-260.

PARADICHLOROBENZENE


fact sheet



23. Kumar, N.; Dale, L. C.; Wijdicks, E. F. M. Mothball mayhem: relapsing toxic leukoencephalopathy due to p-dichlorobenzene neurotoxicity. *Ann. Intern. Med.* 2009, 150 (5), 362-363.
24. Feuillet, L.; Stephanie, M.; Spadari, M. Twin girls with neurocutaneous symptoms caused by mothball intoxication. *N. Engl. J. Med.* 2006, 35 (4), 423-424.
25. Versonnen, B. J.; Arijs, K.; Verslycke, T.; Lema, W.; Janssen, C. R. In vitro and in vivo estrogenicity and toxicity of o-, m-, and p-dichlorobenzene. *Environ. Toxicol. Chem.* 2003, 22 (2), 329-335.
26. NTP Toxicology and Carcinogenesis Studies of 1,4-Dichlorobenzene in F344/N Rats and B6C3F1 Mice; U.S. Department of Health and Human Services, National Institutes of Health. https://ntp-server.niehs.nih.gov/htdocs/LT_rpts/tr319.pdf (accessed Feb 2010), updated Jan 1987.
27. Butterworth, B. E.; Aylward, L. L.; Hays, S. M. A mechanism-based cancer risk assessment for 1,4-dichlorobenzene. *Regul. Toxicol. Pharmacol.* 2007, 49, 138-148.
28. Kokel, D.; Li, Y.; Qin, J.; Xue, D. The nongenotoxic carcinogens naphthalene and para-dichlorobenzene suppress apoptosis in *Caenorhabditis elegans*. *Nat. Chem. Biol.* 2006, 2 (6), 38-345.
29. Aiso, S.; Takeuchi, T.; Arito, H.; Nagano, K.; Yamamoto, S.; Matsushima, T. Carcinogenicity and chronic toxicity in mice and rats exposed by inhalation to para-dichlorobenzene for two years. *J. Vet. Med. Sci.* 2005, 67 (10), 1019-1029.
30. Umemura, I.; Takada, K.; Ogawa, Y.; Kamata, E.; Saito, M.; Kurokawa, Y. Sex difference in inhalation toxicity of p-dichlorobenzene (p-DCB) in rats. *Toxicol. Lett.* 1990, 52, 209-214.
31. Dichlorobenzenes. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans; World Health Organization, International Agency for Research on Cancer: Lyon, France, 1999; Vol. 73, pp 223-265.
32. Neeper-Bradley, T.; Kubena, F. Developmental toxicity study of maternally inhaled paradichlorobenzene vapor in CD rats. Unpublished lab project no. 91N0110, 1992, submitted to U.S. Environmental Protection Agency prepared by Union Carbide Chemicals and Plastics Company, Inc. Revised Reregistration Eligibility Decision (RED) Para-dichlorobenzene, EPA 738-R-07-010; U.S. Environmental Protection Agency, Office of Prevention, Pesticides, and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 2008.
33. Hayes, W.; Gushow, T.; John, J. para-Dichlorobenzene: inhalation teratology study in rabbits. Unpublished report file no. k-1323-(12), 1982, submitted to U.S. Environmental Protection Agency prepared by Dow Chemicals USA. Revised Reregistration Eligibility Decision (RED) Paradichlorobenzene, EPA 738-R-07-010; U.S. Environmental Protection Agency, Office of Prevention, Pesticides, and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 2008.
34. Tyl, R.; Neeper-Bradley, T. Paradichlorobenzene: two-generation reproduction study of inhaled paradichlorobenzene in Sprague-Dawley (CD) rats. Unpublished lab project no. 86-81-90606, 1989, submitted to U.S. Environmental Protection Agency prepared by Union Carbide Bushy Run Research Center. Revised Reregistration Eligibility Decision (RED) Para-dichlorobenzene, EPA 738-R-07-010; U.S. Environmental Protection Agency, Office of Prevention, Pesticides, and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 2008.

PARADICHLOROBENZENE


fact sheet



35. Yoshida, T.; Andoh, K.; Kosaka, H.; Kumagai, S.; Matsunaga, I.; Akasaka, S.; Nakamura, S.; Oda, H.; Fukuhara, M. Inhalation toxicokinetics of p-dichlorobenzene and daily absorption and internal accumulation in chronic low-level exposure to humans. *Arch. Toxicol.* 2002, 76, 306-315.
36. Hill, R. H., Jr.; Ashley, D. L.; Head, S. L.; Needham, L. L.; Pirkle, J. L. p-Dichlorobenzene exposure among 1,000 adults in the United States. *Arch. Environ. Health* 1995, 50 (4), 277-280.
37. Morita, M.; Ohi, G. Para-dichlorobenzene in human tissue and atmosphere in Tokyo metropolitan area. *Environ. Pollut.* 1975, 8, 269-273.
38. Jan, J. Chlorobenzene residues in human fat and milk. *Bull. Environ. Contam. Toxicol.* 1983, 30, 595-599.
39. Hawkins, D. R.; Chasseaud, L. F.; Woodhouse, R. N.; Cresswell, D. G. The distribution, excretion, and biotransformation of p-dichloro[14C]benzene in rats after repeated inhalation, oral and subcutaneous doses. *Xenobiotica* 1980, 10 (2), 81-95.
40. Bomhard, E. M.; Schmidt, U.; Loser, E. Time course of enzyme induction in liver and kidneys and absorption, distribution and elimination of 1,4-dichlorobenzene in rats. *Toxicol.* 1998, 131, 73-91.
41. Hissink, A. M.; Dunnewijk, R.; van Ommen, B.; van Bladeren, P. J. Kinetics and Metabolism of 1,4-Dichlorobenzene in Male Wistar Rats: No Evidence for Quinone Metabolites. *Chem. Biol. Interact.* 1997, 103, 17-33.
42. Klos, C.; Dekant, W. Comparative Metabolism of the Renal Carcinogen 1,4-Dichlorobenzene in Rat: Identification and Quantitation of Novel Metabolites. *Xenobiotica* 1994, 24, 965-976.
43. Ashley, D. L.; Bonin, M. A.; Cardinali, F. L.; McCraw, J. M.; Holler, J. S.; Needham, L. L.; Patterson, D. G., Jr. Determining volatile organic compounds in human blood from a large sample population using purge and trap gas chromatography/mass spectrometry. *Anal. Chem.* 1992, 64, 1021-1029.
44. Hill, R. H., Jr.; Shealy, D. B.; Head, S. L.; Williams, C. C.; Bailey, S. L.; Gregg, M.; Baker, S. E.; Needham, L. L. Determination of pesticide metabolites in human urine using an isotope dilution technique and tandem mass spectrometry. *J. Anal. Toxicol.* 1995, 19, 323-329.
45. Yoshida, T.; Andoh, K.; Fukuhara, M. Urinary 2,5-dichlorophenol as a biological index for p-dichlorobenzene exposure in the general population. *Arch. Environ. Contam. Toxicol.* 2002, 43, 481-485.
46. Fourth national report on human exposure to environmental chemicals; Department of Health and Human Services, Centers for Disease Control and Prevention: Washington, DC, 2009; pp 1, 461-463.
47. Sexton, K.; Adgate, J. L.; Church, T. R.; Ashley, D. L.; Needham, L. L.; Ramachandran, G.; Fredrickson, A. L.; Ryan, A. D. Children's exposure to volatile organic compounds as determined by longitudinal measurements in blood. *Environ. Health Perspect.* 2005, 113 (3), 342-349.
48. Hill, R. H., Jr.; To, T.; Holler, J. S.; Fast, D. M.; Smith, S. J.; Needham, L. L.; Binder, S. Residues of chlorinated phenols and phenoxy acid herbicides in the urine of Arkansas children. *Arch. Environ. Contam. Toxicol.* 1989, 18, 469-474.
49. Wang, M-J.; Jones, K. C. Behavior and Fate of Chlorobenzenes in Spiked and Sewage Sludge-Amended Soil. *Environ. Sci. Technol.* 1994, 28 (11), 1843-1852.

PARADICHLOROBENZENE


fact sheet



50. Spain, J. C.; Nishino, S. F., Degradation of 1,4-dichlorobenzene by a *Pseudomonas* sp. *Appl. Environ. Microbiol.* 1987, 53 (5), 1010-1019.
51. Schwarzenbach, R. P.; Monar-Kubica, E.; Giger, W.; Wakeham, S. G. Distribution, residence time, and fluxes of tetrachloroethylene and 1,4-dichlorobenzene in Lake Zurich, Switzerland. *Environ. Sci. Technol.* 1979, 13 (11), 1367-1373.
52. Howard, P. H. Large Production and Priority Pollutants: 1,4-Dichlorobenzene. *Handbook of Environmental Fate and Exposure Data for Organic Chemicals*; Lewis Publishers: Chelsea, MI, 1989; Vol. 1, pp 250-262.
53. Bouwer, E. J.; McCarty, P. L. Transformations of halogenated organic compounds under denitrification conditions. *Appl. Environ. Microbiol.* 1983, 45 (4), 1295-1299.
54. Bouwer, E. J.; McCarty, P. L. Removal of trace chlorinated organic compounds by activated carbon and fixed-film bacteria. *Environ. Sci. Technol.* 1982, 16 (12), 836-843.
55. Oliver, B. G.; Nicol, K. D. Chlorobenzenes in sediments, water, and selected fish from Lakes Superior, Huron, Erie, and Ontario. *Environ. Sci. Technol.* 1982, 16 (8), 532-536.
56. Schwarzenbach, R. P.; Giger, W.; 56. Hoehn, E.; Schneider, J. K. Behavior of organic compounds during infiltration of river water to ground water. Field studies. *Environ. Sci. Technol.* 1983, 17, 472-479.
57. Young, D. R.; Heesen, T. C. DDT, PCB, and chlorinated benzenes in the marine ecosystem off Southern California. *Water Chlorination: Environmental Impact and Health Effects*; Jolley, R. L.; Hamilton, D. H.; Gorchev, H., Eds.; Ann Arbor Science: Ann Arbor, MI, 1980; pp 267-290.
58. Atkinson, R. Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds under atmospheric conditions. *Chem. Rev.* 1985, 85, 69-201.
59. Ligocki, M. P.; Leuenberger, C.; Pankow, J. F. Trace organic compounds in rain- II. Gas scavenging of neutral organic compounds. *Atmos. Environ.* 1985, 19 (10), 1609-1617.
60. Wang, M.-J.; Jones, K. C. Uptake of Chlorobenzenes by Carrots from Spiked and Sewage Sludge-Amended Soil. *Environ. Sci. Technol.* 1994, 28 (7), 1260-1267.
61. Zhang, J.; Zhao, W.; Pan, J.; Qiu, L.; Zhu, Y. Tissue-dependent distribution and accumulation of chlorobenzenes by vegetables in urban area. *Environ. Int.* 2005, 31, 855-860.
62. Monferran, M. V.; Wunderlin, D. A.; Nimptsch, J.; Pflugmacher, S. Biotransformation and antioxidant response in *Ceratophyllum demersum* experimentally exposed to 1,2- and 1,4-dichlorobenzene. *Chemosphere* 2007, 68, 2073-2079.
63. Brown, S. K.; Sim, M. R.; Abramson, M. J.; Gray, C. N. Concentrations of Volatile Organic Compounds in Indoor Air: A Review. *Indoor Air* 1994, 4 (2), 123-134.
64. Kostianen, R. Volatile organic compounds in the indoor air of normal and sick houses. *Atmos. Environ.* 1995, 29 (6), 693-702.
65. Field, R. A.; Phillips, J. L.; Goldstone, M. E.; Lester, J. N.; Perry, R. Indoor/outdoor interactions during an air pollution event in Central London. *Environ. Technol.* 1992, 13 (4), 391 - 408.
66. Shinohara, N.; Ono, K.; Gamo, M. p-Dichlorobenzene emission rates from moth repellents and leakage rates from cloth storage cases. *Indoor Air* 2008, 18, 63-71.

PARADICHLOROBENZENE

fact sheet



67. De Coensel, N.; Desmet, K.; Sandra, P.; Gorecki, T. Domestic sampling: exposure assessment to moth repellent products using ultrasonic extraction and capillary GC-MS. *Chemosphere* 2008, 71, 711-716.
68. Total Diet Study Market Baskets 1991-3 through 2003-4; U.S. Food and Drug Administration, Center for Food Safety and Applied Nutrition, Office of Food Safety: College Park, MD, 2008; pp i, 51-52.
69. Wang, M-J.; Jones, K. C. Occurrence of chlorobenzenes in nine United Kingdom retail vegetables. *J. Agric. Food Chem.* 1994, 42 (10), 2322-2328.
70. Oliver, B. G.; Nicol, K. D. Chlorobenzenes in sediments, water, and selected fish from Lakes Superior, Huron, Erie, and Ontario. *Environ. Sci. Technol.* 2002, 16 (8), 532-536.
71. Pereira, W. E.; Rostad, C. E.; Chiou, C. T.; Brinton, T. I.; Barber, L. B.; Demcheck, D. K.; Demas, C. R. Contamination of estuarine water, biota, and sediment by halogenated organic compounds: a field study. *Environ. Sci. Technol.* 2002, 22 (7), 772-778.
72. Chung, H. Y. Volatile Components in Crabmeats of *Charybdis feriatus*. *J. Agric. Food Chem.* 1999, 47 (6), 2280-2287.
73. Calamari, D.; Galassi, S.; Setti, F.; Vighi, M. Toxicity of selected chlorobenzenes to aquatic organisms. *Chemosphere* 1983, 12 (2), 253-262.
74. Heitmuller, P. T.; Hollister, T. A.; Parrish, P. R. Acute toxicity of 54 industrial chemicals to sheepshead minnows (*Cyprinodon variegatus*). *Bull. Environ. Contam. Toxicol.* 1981, 27 (1), 596-604.
75. Buccafusco, R. J.; Ells, S. J.; LeBlanc, G. A. Acute toxicity of priority pollutants to bluegill (*Lepomis macrochirus*). *Bull. Environ. Contam. Toxicol.* 1981, 26, 446-452.
76. McPherson, C. A.; Tang, A.; Chapman, P. M.; Taylor, L. A.; Gormican, S. J. Toxicity of 1,4-dichlorobenzene in sediments to juvenile polychaete worms. *Mar. Pollut. Bull.* 2002, 44, 1405-1414.
77. Chapman, P. M.; Paine, M. D.; Arthur, A. D.; Taylor, L. A. A triad study of sediment quality associated with a major, relatively untreated marine sewage discharge. *Mar. Pollut. Bull.* 1996, 32 (1), 47-64.
78. Mortimer, M. R.; Connell, D. W., Critical internal and aqueous lethal concentrations of chlorobenzenes with the crab *Portunus pelagicus* (L). *Ecotoxicol. Environ. Saf.* 1994, 28 (3), 298-312.
79. Mortimer, M. R.; Connell, D. W. Effect of exposure to chlorobenzenes on growth rates of the crab *Portunus pelagicus* (L). *Environ. Sci. Technol.* 1995, 29 (8), 1881-1886.
80. Wong, P. T. S.; Chau, Y. K.; Rhamey, J. S.; Docker, M. Relationship between water solubility of chlorobenzenes and their effects on a freshwater green alga. *Chemosphere* 1984, 13 (9), 991-996.
81. CDC. NIOSH Pocket Guide to Chemical Hazards: p-Dichlorobenzene; U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. <https://www.cdc.gov/niosh/npg/npgd0190.html> (accessed April 2010), updated Feb 2009.
82. ACGIH. TLVs and BEIs, Based on the Documentation of the Threshold Limit Values for Chemical Substances and Physical Agents and Biological Exposure Indices; American Conference of Governmental Hygienists Worldwide: Cincinnati, OH, 2008; pp 24, 42.
83. Basic Information about p-dichlorobenzene in drinking water; U.S. Environmental Protection Agency, Drinking Water Contaminants. <https://www.epa.gov/safewater/contaminants/basicinformation/p-dichlorobenzene.html> (accessed April 2010), updated March 2010.