
fact sheet



What is naphthalene?

Naphthalene is made from crude oil or coal tar. It is also produced when things burn, so naphthalene is found in cigarette smoke, car exhaust, and smoke from forest fires. It is used as an <u>insecticide</u> and pest repellent. Naphthalene was first registered as a pesticide in the United States in 1948.

What are some products that contain naphthalene?

Mothballs and other products containing naphthalene are solids that turn into toxic gas. The toxic gas kills insects like <u>pantry moths</u> and may repel animals. There are over a dozen products containing naphthalene registered for use by the U.S. Environmental Protection Agency (U.S. EPA).



Photo credit: NPIC

How does naphthalene work?

When naphthalene gas is inhaled, the body breaks it down into other chemicals that react with cells in the body and damage tissues. How naphthalene kills <u>moths</u> is not understood. The odor is thought to be unpleasant enough to drive animals away in repellent products.

How might I be exposed to naphthalene?


You can be exposed to a pesticide if you breathe it in, get it on your skin, or if you accidentally eat or drink something containing a pesticide. This can happen if you get some on your hands and don't wash them before eating or smoking. People are most likely to be exposed to naphthalene by breathing in the vapors. When you smell mothballs, you are inhaling the pesticide. Small children and pets are at risk of eating mothballs, because they look like candy or other treats.

IMPORANT: Always <u>follow label instructions</u> and take steps to <u>avoid exposure</u>. If any exposures occur, be sure to follow the First Aid instructions on the product label carefully. For additional treatment advice, contact the Poison Control Center at 800-222-1222. If you wish to discuss a <u>pesticide problem</u>, please call 1-800-858-7378.

What are some signs and symptoms from a brief exposure to naphthalene?

People have developed headaches, nausea, dizziness, and/or vomiting after being exposed to naphthalene vapors. If someone breathes in enough of the vapor or eats a mothball containing naphthalene, they might develop hemolytic anemia. This is when red blood cells break apart, and no longer carry oxygen the way they should. Small children have also developed diarrhea, fever, abdominal pain, and painful urination with discolored urine after eating naphthalene mothballs. Dogs that have eaten naphthalene mothballs may have lethargy, vomiting, diarrhea, lack of appetite and tremors.

fact sheet



Clothing that was stored in mothballs without being washed afterwards has caused anemia in infants who wore the clothing, diapers or blankets. People with an inherited enzyme deficiency are at much greater risk of anemia than people with normal enzyme levels.

What happens to naphthalene when it enters the body?

In humans, naphthalene is broken down to alpha-naphthol, which is linked to the development of hemolytic anemia. Kidney and liver damage may also occur. Alpha-naphthol and other metabolites are excreted in urine.

In animals, naphthalene breaks down into other compounds including alpha-naphthol, which may affect the lungs and eyes. Naphthalene was found in the milk of exposed cows, but the residues disappeared quickly after the cows were no longer exposed. Nearly all the naphthalene was broken down into other compounds and excreted in their urine.

Is naphthalene likely to contribute to the development of cancer?

Animal studies have suggested that naphthalene can cause cancer. The International Agency for Research on Cancer (IARC) of the World Health Organization (WHO) concluded that naphthalene is possibly carcinogenic to humans. The U.S. EPA classified naphthalene as a possible human carcinogen, also based on animal studies.

Has anyone studied non-cancer effects from long-term exposure to naphthalene?

Rats fed naphthalene while pregnant did not gain weight as quickly. In humans, women who ate naphthalene mothballs or inhaled the vapors while pregnant gave birth to babies with hemolytic anemia. No information was found on naphthalene and asthma or other chronic diseases.


Are children more sensitive to naphthalene than adults?

While <u>children</u> may be especially sensitive to pesticides compared to adults, there are currently no data showing that children have increased sensitivity specifically to naphthalene. However, small children are at greatest danger from eating stray mothballs, because they may look like candy.

What happens to naphthalene in the environment?

Most naphthalene in the environment will turn into a gas. Some of it may be bound to soil, where it can be taken up by plants. It can also be deposited on plant leaves from the air. Naphthalene is broken down by bacteria, fungi, air, and sunlight. Naphthalene has been found in wastewater treatment plant discharge. No information was found on naphthalene and groundwater. The half-life of naphthalene in the environment may range from less than one day in air to over 80 days in soil.

fact sheet



Can naphthalene affect birds, fish, or other wildlife?

Naphthalene was considered moderately toxic to several species of fish, water fleas, and Pacific oysters. It was considered slightly toxic to green algae. Naphthalene was considered practically non-toxic after being fed to bobwhite quail.

Where can I get more information?


For more detailed information about naphthalene please visit the list of referenced resources below, call the National Pesticide Information Center, Monday - Friday, between 8:00am - 12:00pm Pacific Time (11:00am - 3:00pm Eastern Time) at 1-800-858-7378 or visit us on the web at npic.orst.edu. NPIC provides objective, science-based answers to questions about pesticides.

Date Reviewed: December 2010


Please cite as: Gervais, J.; Luukinen, B.; Buhl, K.; Stone, D. 2010. *Naphthalene General Fact Sheet*; National Pesticide Information Center, Oregon State University Extension Services. http://npic.orst.edu/factsheets/naphgen.html.

References:


- 1. *Toxicological Profile for Naphthalene, 1-methylnaphthalene, and 2-methylnaphthalene*; U.S Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry: Atlanta, GA, 2010.
- 2. Bischoff, K. Naphthalene. *Clinical Veterinary Toxicology*; Plumlee, K. H., Ed.; Mosby: St. Louis, MO, 2004, pp. 163-164.
- 3. Reregistration Eligibility Decision (RED): Naphthalene; EPA 738-R-07-010; U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 2008.
- 4. WHO. *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*; International Agency for Research on Cancer, World Health Organization: Lyon, France, 2002; pp 367-435.
- 5. Preuss, R.; Angerer, J.; Drexler, H. Naphthalene An environmental and occupational toxicant. *Int. Arch. Occup. Environ. Health* 2003, 76, 556-576.
- 6. *Hazardous Substances Data Bank (HSDB): Naphthalene*; U.S. Department of Health and Human Services, National Institutes of Health, National Library of Medicine: Bethesda, MD, 2005.
- 7. WHO. *Naphthalene Poisons Information Monograph 363 (PIM 363)*; International Programme on Chemical Safety, World Health Organization: Lyon, France, 2000.
- 8. Bogen, K. T.; Benson, J. M.; Yost, G. S.; Morris, J. B.; Dahl, A. R.; Clewell, H. J., III; Krishnan, K.; Omiecinski, C. J. Naphthalene metabolism in relation to target tissue anatomy, physiology, cytotoxicity and tumorigenic mechanism of action. *Reg. Toxicol. Pharmacol.* 2008, 51, S27-S36.




- 9. Buckpitt, A. R.; Boland, B.; Isbell, M.; Morin, D.; Shultz, M.; Baldwin, R.; Chan, K.; Karlsson, A.; Lin, C.; Taff, A.; West, J.; Fanucchi, M.; Van Winkle, L.; Plopper, C. G. Naphthalene-induced respiratory tract toxicity: metabolic mechanisms of toxicity. *Drug Metab. Rev.* 2002, 34 (4), 791-820.
- 10. Reigart, J. R.; Roberts, J. R. Fumigants. *Recognition and Management of Pesticide Poisonings, 5th ed.*; U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 1999; pp 156-162.
- 11. Valaes, T.; Doxiadis, S. A.; Fessas, P. Acute hemolysis due to naphthalene inhalation. *J. Pediatr.* 1963, 63, 904-915.
- 12. Santucci, K.; Shah, B. R. Association of naphthalene with acute hemoyltic anemia. *Acad. Emerg. Med.* 2000, 7, 42-47.
- 13. Molloy, E. J.; Doctor, B. A.; Reed, M. D.; Walsh, M. C. Perinatal toxicity of domestic naphthalene exposure. *J. Perinatol.* 2004, 24, 792-793.
- 14. Phimister, A. J.; Lee, M. G.; Morin, D.; Buckpitt, A. R.; Plopper, C. G. Glutathione depletion is a major determinant of inhaled naphthalene respiratory toxicity and naphthalene metabolism in mice. *Toxicol. Sci.* 2004, 82, 268-278.
- 15. Phimister, A. J.; Nagasawa, H. T.; Buckpitt, A. R.; Plopper, C. G. Prevention of naphthalene-induced pulmonary toxicity by glutathione prodrugs: roles for glutathione depletion in adduct formation and cell injury. *J. Biochem. Mol. Toxicol.* 2005, 19 (1), 42-51.
- 16. Wilson, A. S.; Davis, C. D.; Williams, D. P.; Buckpitt, A. R.; Pirmohamed, M.; Park, B. K. Characterization of the toxic metabolite(s) of naphthalene. *Toxicol.* 1996, 114, 233-242.
- 17. Papciak, R. J.; Malloy, V. T. Acute toxicological evaluation of naphthalene. *J. Am. Coll. Toxicol.* 1990, 1 (1), 17-19.
- 18. Gaines, T. B. Acute toxicity of pesticides. *Toxicol. Appl. Pharmacol.* 1969, 14, 515-534.
- 19. Shopp, G. M.; White, K. L., Jr.; Holsapple, M. P.; Barnes, D. W.; Duke, S. S.; Anderson, A. C.; Condie, L. W.; Hayes, J. R.; Borzelleca, J. F. Naphthalene toxicity in CD-1 mice: general toxicology and immunotoxicology. *Fundam. Appl. Toxicol.* 1984, 4, 406-419.
- 20. Plasterer, M. R.; Bradshaw, W. S.; Booth, G. M.; Carter, M. W. Developmental toxicity of nine selected compounds following prenatal exposure in the mouse: naphthalene, p-nitrophenol, sodium selenate, dimethyl phthalate, ethylenethiourea, and four glycol ether derivatives. *J. Toxicol. Environ. Health* 1985, 15, 25-38.
- 21. West, J. A. A.; Pakehham, G.; Morin, D.; Fleschner, C. A.; Buckpitt, A. R.; Plopper, C. G., Inhaled naphthalene causes dose dependent clara cell cytotoxicity in mice but not in rats. *Toxicol. Appl. Pharmacol.* 2001, 173, 14-119.
- 22. Witschi, H. R.; Pinkerton, K. E.; Van Winkle, L. S.; Last, J. A. Toxic responses of the respiratory system. *Casarette and Doull's Toxicology: The basic science of poisons, 7th ed.*; Klaassen, C. D., Ed.; McGraw Hill Medical: New York, 2008; p 613.
- 23. Zuelzer, W. W.; Apt, L. Acute hemolytic anemia due to naphthalene poisoning. *J. Am. Med. Assoc.* 1949, 141 (3), 185-190.



- 24. Desnoyers, M.; Hebert, P. Heinz body anemia in a dog following possible naphthalene ingestion. *Vet. Clin. Pathol.* 1995, 24 (4), 124-125.
- 25. Health Effects Support Document for Naphthalene; EPA 822-R-03-005; U.S. Environmental Protection Agency, Office of Water, Health and Ecological Criteria Division, U.S. Governement Printing Office: Washington, DC, 2003.
- 26. Grant, W. M. Toxicology of the Eye, 3rd ed.; Charles C. Thomas: Springfield, IL, 1986, pp 650-645.
- 27. Coombs, D. 27. Naphthalene 4-week inhalation study in rats. Unpublished lab project no. LDA 1/921559, 1993, submitted to U.S. Environmental Protection Agency by Huntingdon Research Center, Ltd. EPA MRID 42934901. *Reregistration Elegibility Decision (RED) Naphthalene*; U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 2008, p 14.
- 28. Frantz, S.; Van Miller, J.; Papciak, R. Ninety-day (sub-chronic) dermal toxicity study with naphthalene in albino rats: final report. Unpublished project no. 49- 539, 1986, submitted to U.S. Environmental Protection Agency by Union Carbide Bushy Run Research Center. EPA MRID 40021801. *Reregistration Elegibility Decision (RED) Naphthalene*; U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 2008, p 14.
- 29. Shah, B. R.; Santucci, K. Naphthalene induced acute hemolytic-anemia in children with glucose-6-phosphate-dehydrogenase (G-6-PD) deficiency naphthalene has no legitimate place on the market as a moth repellent. *Pediatr. Res.* 1995, 37 (4), A144.
- 30. Harden, R. A.; Baetjer, A. M. Aplastic anemia following exposure to paradichlorobenzene and naphthalene. *J. Occup. Med.* 1978, 20 (12), 820-822.
- 31. Thomas, P.; Budiantara, L. Reproductive life history stages sensitive to oil and naphthalene in Atlantic croaker. *Mar. Environ. Res.* 1995, 39, 147-150.
- 32. Abdo, K. M.; Grumbein, S.; Chou, B. J.; Herbert, R. Toxicity and carcinogenicity study in F344 rats following 2 years of whole-body exposure to naphthalene vapors. *Inhal. Toxicol.* 2001, 13, 931-950.
- 33. NTP Technical Report on the toxicology and carcinogenesis studies of naphthalene (CAS No. 91-20-3) in B6C3F1 mice; U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, National Toxicology Program: Research Triangle Park, NC, 1992; pp 35-36.
- 34. Kokel, D.; Li, Y.; Qin, J.; Xue, D., The nongenotoxic carcinongens naphthalene and paradichlorobenzene suppress apoptosis in Caenorhabditis elegans. *Nat. Chem. Biol.* 2006, 2 (6), 338-345.
- 35. Integrated Risk Information System (IRIS): Naphthalene (CASRN 91-20-3); U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC, 2008.
- 36. Developmental toxicity study in rabbits: naphthalene. Unpublished report number PH 329-TX-001 85, 1986, submitted to U.S. Department of Health and Human Services by Pharmakon Research International, Inc. *Toxicologial Profile for naphthalene, 1-methylnaphthalene, and 2-methylnaphthalene*; U.S Department of Health and Human Services, Public Health Service, Agency for Toxic Subtances and Disease Registry: Atlanta, 2005; p 75.




- 37. Developmental toxicity of naphthalene (CAS No. 91-20-3) administered by gavage to Sprague-Dawley (CD) rats on gestational days 6 through 15: Final study report and appendix; U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, National Toxicology Program, TER91006: Research Triangle Park, NC, 1991, p 76.
- 38. Anziulewicz, J. A.; Dick, H. J.; Chiarulli, E. E. Transplacental naphthalene poisoning. *Am. J. Obstet. Gynecol.* 1959, 78 (3), 519-521.
- 39. Pellizzari, E. D.; Hartwell, T. D.; Harris, B. S. H., III; Waddell, R. D.; Whitaker, D. A.; Erickson, M. D. Pureable organic compounds in mother's milk. *Bull. Environ. Contam. Toxicol.* 1982, 28, 322-328.
- 40. Lehmann, I.; Thoelke, A.; Rehwagen, M.; Rolle-Kampczyk; Schlink, U.; Schulz, R.; Borte, M.; Diez, U.; Herbarth, O. The influence of maternal exposure to volatile organic compounds on the cytokine secretion profile of neonatal T cells. *Environ. Toxicol.* 2002, 17 (3), 203-210.
- 41. Turkall, R. M.; Skowronsi, G. A.; Abdel-Rahman, M. S. A comparative study of the kinetics and bioavailability of pure and soil-adsorbed naphthalene in dermally exposed male rats. *Arch. Environ. Contam. Toxicol.* 1994, 26, 504-509.
- 42. Schafer, W. B. Acute hemolytic anemia related to naphthalene. *Pediatrics* 1951, 7, 172-174.
- 43. Buckpitt, A. R., Evidence for hepatic formation, export and covalent binding of reactive naphthalene metabolites in extrahepatic tissues in vivo. J. Pharmacol. Exp. Ther. 1983, 225 (1), 8-16. 44. Eisele, G. R. Naphthalene distribution in tissues of laying pullets, swine, and dairy cattle. *Bull. Environ. Contam. Toxicol.* 1985, 34, 549-556.
- 44. O'Brien, K. A. F.; Suverkropp, C.; Kanekal, S.; Plopper, C. G.; Buckpitt, A. R. Tolerance to multiple doses of the pulmonary toxicant, naphthalene. *Toxicol. App. Pharmacol.* 1989, 99, 487-500.
- 45. West, J. A. A.; Buckpitt, A. R.; Plopper, C. G. Elevated air 46. way GSH reysnthesis confers protection to Clara cells from naphthalene injury in mice made tolerant by repeated exposures. *J. Pharmacol. Exp.* Ther. 2000, 294 (2), 516-523.
- 46. Van Heyningen, R.; Pirie, A. The metabolism of naphthalene and its toxic effect on the eye. *Biochem. J.* 1967, 102, 842-852.
- 47. Bieniek, G. Urinary naphthols as an indicator of exposure to naphthalene. Scand. *J. Work Environ. Health* 1997, 23, 414-420.
- 48. Bieniek, G., The presence of 1-naphthol in the urine of industrial workers exposed to naphthalene. *Occup. Environ. Med.* 1994, 51, 357-359.
- 49. Yang, M.; Koga, M.; Tawamoto, T. A study for the proper application of urinary naphthols, new biomarkers for airborne polycyclic aromatic hydrocarbons. *Arch. Environ. Contam. Toxicol.* 1999, 36, 99-108.
- 50. Howard, P. H. Naphthalene. *Handbook of Environmental Fate and Exposure Data for Organic Chemcials*; Jarvis, W. F.; Sage, G. W.; Basu, D. K.; Gray, D. A.; Meylan, W.; Crosbie, E. K., Eds; Lewis Publishers: Chelsea, MI, 1989; Vol. 1, pp 408-422.
- 51. Cerniglia, C. E. Biodegradation of polycyclic aromatic hydrocarbons. *Biodegradation* 1992, 3, 351-368.



- 52. Jury, W. A.; Spencer, W. F.; Farmer, W. J. Behavior assessment model for trace organics in soil: III. Application of screening model. *J. Environ. Qual.* 1984, 13 (4), 573-479.
- 53. Treccani, V.; Walker, N.; Wiltshire, G. H., The metabolism of naphthalene by soil bacteria. *J. Gen. Microbiol.* 1954, 11, 341-348.
- 54. Cerniglia, C. E., Microbial metabolism of polycyclic aromatic hydrocarbons. *Adv. Appl. Microbiol.* 1984, 30, 31-71.
- 55. Bouwer, E. J.; McCarty, P. L.; Bouwer, H.; Rice, R. C. Organic contaminant behavior during rapid infiltration of secondary wastewater at the Phoenix 23rd Avenue Project. *Water Res.* 1984, 18 (4), 463-472.
- 56. Southworth, G. R. The role of volatilization in removing polycyclic aromatic hydrocarbons from aquatic environments. *Bull. Environ. Contam. Toxicol.* 1979, 21, 507-514.
- 57. Zepp, R. G.; Schlotzhauer, P. F. Influence of algae on photolysis rates of chemicals in water. *Environ. Sci. Technol.* 1983, 17, 462-466.
- 58. Cerniglia, C. E.; van Baalen, C.; Gibson, D. T., Metabolism of naphthalene by the cyanobacterium Oscillatoria sp., strain JCM. *J. Gen. Microbiol.* 1980, 116, 485-494.
- 59. Cerniglia, C. E.; Gibson, D. T.; van Baalen, C. Oxidation of naphthalene by cyanobacteria and microalgae. *J. Gen. Microbiol.* 1980, 116, 495-500.
- 60. Atkinson, R. Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds under atmospheric conditions. *Chem. Rev.* 1986, 86, 69-201.
- 61. Atkinson, R.; Arey, J.; Zielinska, B.; Aschmann, S. M., Kinetics and products of the gas-phase reactions of OH radicals and N2O5 with naphthalene and biphenyl. *Environ. Sci. Technol.* 1987, 21, 1014-1022.
- 62. Gao, Y.; Collins, C. D. Uptake pathways of polycyclic aromatic hydrocarbons in white clover. *Environ. Sci. Technol.* 2009, 43 (16), 6190-6195.
- 63. Schwab, A. P.; Al-Assi, A. A.; Banks, M. K. Adsorption of naphthalene onto plant roots. *J. Environ. Qual.* 1998, 27, 220-224.
- 64. Watts, A. W.; Ballestero, T. P.; Gardner, K. H. Uptake of polycyclic aromatic hydrocarbons (PAHs) in salt marsh plants Spartina alterniflora grown in contaminated sediments. *Chemosphere* 2006, 62, 1253-1260.
- 65. Kipopoulou, A. M.; Manoli, E.; Samara, C. Bioconcentration of polycyclic aromatic hydrocarbons in vegetables grown in an industrial area. *Environ. Pollut.* 1999, 106, 369-380.
- 66. De Coensel, N.; Desmet, K.; Sandra, P.; Gorecki, T. Domestic sampling: exposure assessment to moth repellent products using ultrasonic extraction and capillary GC-MS. *Chemosphere* 2008, 71, 711-716.
- 67. Chuang, J. C.; Callahan, P. J.; Lyu, C. W.; Wilson, N. K. Polycyclic aromatic hydrocarbon exposures of children in low-income families. *J. Expo. Anal. Environ. Epidemiol.* 1999, 2, 85-98.
- 68. Revised Ecological Risk Assessment for the Reregistration Elegibility Decision (RED) for Naphthalene; U.S. Environmental Protection Agency, Offices of Prevention, Pesticides, and Toxic Substances, Environmental Fate and Effects Division, U.S. Government Printing Office: Washington, DC, 2008.

fact sheet



- 69. Varanasi, U.; Uhler, M.; Stranahan, S. I. Uptake and release of naphthalene and its metabolites in skin and epidermal mucus of salmonids. *Toxicol. Appl. Pharmacol.* 1978, 44, 277-289.
- 70. Kelsey, J. W.; Alexander, M., Declining bioavailability and inappropriate estimation of risk of persistent compounds. *Environ. Toxicol. Chem.* 1997, 16 (3), 582-585.
- 71. *NIOSH Pocket Guide to Chemical Hazards: Naphthalene*; U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institure for Occupational Safety and Health: Atlanta, GA, 2005.

NPIC es un acuerdo cooperativo entre la Universidad Estatal de Oregon y la Agencia de Protección Ambiental de EE.UU. (US EPA). La información contenida en esta publicación de ninguna manera reemplaza o supersede las restricciones, precauciones, direcciones u otra información en la etiqueta del plaguicida o cualquier otro requisito reglamentario, ni refleja necesariamente la posición de la EPA de Estados Unidos.

